

Lecture #20: Random Variables and Expectation

Key Concepts

Some — but not all — of this material is review material.

Random Variables

Definition 1. Let Ω be a sample space. A **random variable over Ω** is a (total) function $X : \Omega \rightarrow \mathbb{R}$.

We will often be interested in random variable whose ranges are particular *subsets* V of $\mathbb{R}$ — so that these functions can also be viewed as functions $X : \Omega \rightarrow V$ (as well as functions $X : \Omega \rightarrow \mathbb{R}$).

Definition 2. An **indicator random variable** over a sample space Ω , is a random variable X such that $X(\sigma) \in \{0, 1\}$ for all $\sigma \in \Omega$.

Once again, let Ω be a sample space and let $X : \Omega \rightarrow \mathbb{R}$.

- We will write “ $X = r$ ” as the name of the event

$$\{\sigma \in \Omega \mid X(\sigma) = r\} \subseteq \Omega.$$

- We will write “ $X \geq r$ ” as the name of the event

$$\{\sigma \in \Omega \mid X(\sigma) \geq r\} \subseteq \Omega.$$

- “ $X \leq r$ ”, “ $X > r$ ”, “ $X < r$ ”, and “ $X \neq r$ ” can be used as the names for (corresponding) events in the same way.

The Expected Value of a Random Variable

Definition 3. Let Ω be a sample space with probability distribution $P : \Omega \rightarrow \mathbb{R}$, and let $X : \Omega \rightarrow \mathbb{R}$ be a random variable over Ω .

Suppose that

$$\sum_{\sigma \in \Omega} P(\sigma) \times |X(\sigma)|$$

is finite — that is, “less than $+\infty$ ”.¹

Then the **expected value of X , with respect to probability distribution P** , is the value

$$E[X] = \sum_{\sigma \in \Omega} P(\sigma) \times X(\sigma).$$

Recall, from the lecture on “Conditional Probability”, that if Ω is a sample space, $P : \Omega \rightarrow \mathbb{R}$ is a probability distribution, and $B \subseteq \Omega$ is an event such that $P(B) > 0$, then a **conditional probability distribution** $P_B : \Omega \rightarrow \mathbb{R}$ can be defined by setting

$$P_B(\sigma) = \begin{cases} \frac{P(\sigma)}{P(B)} & \text{if } \sigma \in B, \\ 0 & \text{if } \sigma \notin B \end{cases}$$

for every outcome $\sigma \in \Omega$.

Definition 4. If X is a random variable then the **conditional expectation of X given B** is the expected value of X with the respect to the conditional probability P_B :

$$E[X | B] = \sum_{\sigma \in \Omega} P_B(\sigma) \times X(\sigma).$$

Linearity of Expectation

Let Ω be a sample space with a probability distribution $P : \Omega \rightarrow \mathbb{R}$, and let $X_1, X_2, \dots, X_n : \Omega \rightarrow \mathbb{R}$ be random variables, for some positive integer n . “ $X_1 + X_2 + \dots + X_n$ ” denotes a random variable such that

$$(X_1 + X_2 + \dots + X_n)(\sigma) = X_1(\sigma) + X_2(\sigma) + \dots + X_n(\sigma)$$

for each outcome $\sigma \in \Omega$. Similarly, if $X : \Omega \rightarrow \mathbb{R}$ and $a, b \in \mathbb{R}$ then “ $a \cdot X + b$ ” denotes a random variable such that

$$(aX + b)(\sigma) = a \cdot X(\sigma) + b$$

for every outcome $\sigma \in \Omega$, as well.

¹This is a “technical restriction” that you will not need to worry about whenever Ω is a finite set.

Claim 1 (Linearity of Expectation). Let Ω be a sample space with probability distribution $P : \Omega \rightarrow \mathbb{R}$.

(a) If $X_1, X_2, \dots, X_n : \Omega \rightarrow \mathbb{R}$ are random variables over Ω , for a positive integer n , then

$$E[X_1 + X_2 + \dots + X_n] = E[X_1] + E[X_2] + \dots + E[X_n].$$

(b) If $X : \Omega \rightarrow \mathbb{R}$ is a random variable over Ω and $a, b \in \mathbb{R}$ then

$$E[a \cdot X + b] = a \cdot E[X] + b.$$

Independent Random Variables

Once again, suppose that Ω is a sample space with probability distribution $P : \Omega \rightarrow \mathbb{R}$. Consider random variables $X : \Omega \rightarrow V_X$ and $Y : \Omega \rightarrow V_Y$, where $V_X, V_Y \subseteq \mathbb{R}$. Recall that, for $a \in V_X$ and $b \in V_Y$,

$$“X = a” = \{\sigma \in \Omega \mid X(\sigma) = a\}$$

and

$$“Y = b” = \{\tau \in \Omega \mid Y(\tau) = b\},$$

so that

$$“X = a \wedge Y = b” = \{\mu \in \Omega \mid X(\mu) = a \text{ and } Y(\mu) = b\}.$$

Definition 5. The above random variables X and Y are **independent** if

$$P(X = a \wedge Y = b) = P(X = a) \times P(Y = b)$$

for all values $a \in V_X$ and $b \in V_Y$.

Claim 2. Let Ω be a sample space with probability distribution $P : \Omega \rightarrow \mathbb{R}$, and let $X, Y : \Omega \rightarrow \mathbb{R}$. If X and Y are independent random variables then

$$E[X \times Y] = E[X] \times E[Y].$$

Pairwise Independent Random Variables

Once again, let Ω be a sample space with probability distribution $P : \Omega \rightarrow \mathbb{R}$. Let k be a positive integer and let

$$X_1 : \Omega \rightarrow V_1, X_2 : \Omega \rightarrow V_2, \dots, X_k : \Omega \rightarrow V_k$$

be random variables over Ω (so that $V_1, V_2, \dots, V_k \subseteq \mathbb{R}$).

Definition 6. The random variables $X_1, X_2, \dots, X_k$ are **pairwise independent** if X_i and X_j are independent, for every pair of numbers i and j such that $1 \leq i < j \leq k$.

Mutually Independent Random Variables

Finally, let Ω be a sample space with probability distribution $P : \Omega \rightarrow \mathbb{R}$. Let k be a positive integer and let

$$X_1 : \Omega \rightarrow V_1, X_2 : \Omega \rightarrow V_2, \dots, X_k : \Omega \rightarrow V_k$$

be random variables over Ω (so that $V_1, V_2, \dots, V_k \subseteq \mathbb{R}$).

Definition 7. The random variables $X_1, X_2, \dots, X_k$ are ***mutually independent*** if the following condition is satisfied: ***For every*** subset $S \subseteq \{1, 2, \dots, k\}$ and ***for all*** combinations of $a_i \in V_i$, for $i \in S$,

$$P\left(\bigwedge_{i \in S} (X_i = a_i)\right) = \prod_{i \in S} P(X_i = a_i).$$